

The original citation of Jerdon's Nightjar *Caprimulgus atripennis* (Caprimulgidae)

NIGEL CLEERE

Thomas Claverhill Jerdon named *Caprimulgus atripennis* in his 'Illustrations of Indian ornithology', which were a collection of descriptions and colour plates of Indian birds published between 1843 and 1847 (Jerdon 1843-1847, Zimmer 1926). The citation of Jerdon's Nightjar in this work has simply been given as plate 24 by all subsequent authors and the publication date for this plate was determined as 1845 by Sherborn (1922).

An examination of this work reveals that plate 24 clearly shows a different species entirely, the Grey Nightjar *Caprimulgus indicus*, and the accompanying text is headed *Caprimulgus indicus*, although with the different English name of Large Indian Nightjar. The identification of the species on plate 24 as *Caprimulgus indicus* can be accurately determined by the greyish ground colour, the lack of a hindneck collar, the bold, buff spots on otherwise uniform wing-coverts and the broad white tips to the four outer tail feathers. *Caprimulgus atripennis* can be ruled out by the lack of rufous-tinged ear-coverts and lores, the lack of a broad rufous hindneck collar and the lack of a bold scapular pattern.

The text commences with a reference to the original description of *Caprimulgus indicus* Latham, 1790 and confirms that identification was supported by reference to Gray (1832), although the illustration (plate 34, fig. 1) in that work is of an utterly nondescript

species. Much of the next two pages of text describe the range, habitat and habits of *Caprimulgus indicus*, although the description of the calls seems to suggest *Caprimulgus atripennis*. On the third page, the author writes about other species of Indian nightjars and it is only under the Large-tailed Nightjar *Caprimulgus macrourus* [sic], that he refers to specimens taken from the eastern Ghauts [Ghats] and proposes the name *Caprimulgus atripennis* for them. Only the length, wing and tail measurements are then given.

As a result of the above examination, I suggest that a more accurate citation for Jerdon's Nightjar is:

Caprimulgus atripennis Jerdon 1845, *Ill. Ind. Orn.*, p. 3 of text to pl. 24.

REFERENCES

- Gray, J. E. (1832) *Illustrations of Indian zoology* Vol. 1. London: Adolphus Richter.
 Jerdon, T. C. (1843-1847) *Illustrations of Indian ornithology*. Madras: American Mission Press.
 Latham, J. (1790) *Index ornithologicus* Vol. 2. London: Leigh and Sotheby.
 Sherborn, C. D. (1922) *Index animalium*. London: British Museum.
 Zimmer, J. T. (1926) Catalogue of the Edward E. Ayer ornithological library. *Field Mus. Nat. Hist.* 16.

Nigel Cleere, 2 Hawthorn House, Roundfield, Upper Bucklebury, Berks. RG7 6RQ, U.K. Email: cleere@churr.freereserve.co.uk

Ashy Wood Pigeon *Columba pulchricollis*: a new species for Vietnam

ANDREW W. TORDOFF

The following observation was made during a 12-month survey of Hoang Lien Nature Reserve, Vietnam, by the environmental conservation NGO, Frontier-Vietnam. Hoang Lien Nature Reserve includes Vietnam's highest mountain, Fan Si Pan (3,143 m).

On 20 August 1998 I was passing through an area of burnt scrub on Fan Si Pan mountain (22°19'N 103°48'E, 1,900 m) when I observed a pigeon land in a burnt tree 20 m away. I had an unobstructed view of the bird perched prominently for approximately one minute, after which it flew to a more distant tree and then out of sight. I immediately noticed the buffy collar with black markings, which contrasted sharply with the dark slate-grey upperparts and grey upper breast.

Neither the upperparts nor the buffy-cream belly had any visible markings. The head was pale grey and the feet were red. I considered this bird to be Ashy Wood Pigeon *Columba pulchricollis*, ruling out the two other potential species after consultation with the relevant literature: (1) Speckled Wood Pigeon *Columba hodgsonii* was ruled out on the basis of the presence in my bird of a buffy collar, the colour of the feet being red rather than grey, and the absence of white speckling on the wing-coverts and belly; (2) Mountain Imperial Pigeon *Ducula badia* was ruled out on the basis of size, the presence of a buffy collar, and the absence of banding in the tail.

Ashy Wood Pigeon is not listed for Vietnam by King *et al.* (1975), Vo Quy and Nguyen Cu (1995), del Hoyo *et al.* (1997), Robson (2000), Gibbs *et al.* (2001) or in any other literature available to the author. I therefore consider this observation to be a new record for Vietnam. Ashy Wood Pigeon occurs between 1,600 and 2,500 m on Doi Inthanon mountain in Thailand (Round 1988). The species is described as 'fairly common' on Mount Victoria, Burma (Robson *et al.* 1998), and there is a single record from northern Laos (Duckworth *et al.* 1999). The species is also known from Yunnan (China), Taiwan, Bhutan, India and Nepal (del Hoyo *et al.* 1997, Gibbs *et al.* 2001). It is not surprising, therefore, that it should be found in a high-altitude habitat in northern Vietnam. The habitat for Ashy Wood Pigeon has been described as hill evergreen forest and secondary growth (Lekagul and Round 1991), and dense mixed evergreen and deciduous hill forest and secondary growth from 1,100 to 3,200 m (Gibbs *et al.* 2001). The occurrence of this species in scrub is not necessarily anomalous as the bird was observed on a burnt ridge-top that was surrounded on all sides by montane evergreen forest.

Fan Si Pan mountain is a relatively well-studied locality in Vietnam (Delacour 1930, Bangs and van Tyne 1931, Korzun and Kalyakin 1998) so it is perhaps surprising that Ashy Wood Pigeon has not been recorded by previous surveys. However, it is known to wander in response to available food supply (J. C. Eames *in litt.*, Gibbs *et al.* 2001), and it may be that this species is only an infrequent visitor to the area. Alternatively, it may be that the high hunting pressure prevalent in Hoang Lien Nature Reserve has significantly reduced the population of this species. It is to be hoped that future conservation efforts can preserve this newly recorded species and the high avian diversity of this area (Tordoff *et al.* 1999).

This work was carried out during the Frontier-Vietnam Forest Research Programme, which is a collaboration between the Society for Environmental Exploration and the Institute of Ecology and Bio-

logical Resources, Hanoi. The author would like to thank Dr Damon Stanwell-Smith of the Society for Environmental Exploration and Jonathan C. Eames of the BirdLife International Vietnam Programme for their comments on the first draft of this paper.

REFERENCES

- Bangs, O. and van Tyne, J. (1931) Birds of the Kelley-Roosevelts expedition to French Indo-china. *Publ. Field Mus. Nat. Hist. (Zool. Ser.)* 18: 33-119.
- del Hoyo, J., Elliott, A. and Sargatal, J., eds. (1997) *Handbook of birds of the world. Vol. 4: sandgrouse to cuckoos*. Barcelona: Lynx Edicions.
- Delacour, J. (1930) On the birds collected during the fifth expedition to French Indochina. *Ibis* (12)6: 564-599.
- Duckworth, J. W., Salter, R. E. and Khounboline, K., compilers (1999) *Wildlife in Lao P.D.R.: 1999 status report*. Vientiane: IUCN, Wildlife Conservation Society and Centre for Protected Areas and Watershed Management.
- Gibbs, D., Barnes, E. and Cox, J. (2001) *Pigeons and doves: a guide to the pigeons and doves of the world*. Mountfield, Sussex, U.K.: Pica Press.
- King, B. F., Dickinson, E. C. and Woodcock, M. W. (1975) *A field guide to the birds of South-East Asia*. London: Collins.
- Korzun, L. P. and Kalyakin, M. V. (1998) [*Materials of zoological and botanical studies in Fan Si Pan summit area (north Vietnam)*]. Moscow and Hanoi: Vietnam-Russia Tropical Centre. (In Russian.)
- Lekagul, B. and Round, P. D. (1991) *A guide to the birds of Thailand*. Bangkok: Saha Karn Bhaet.
- Robson, C. R. (2000) *A field guide to the birds of Thailand and South-East Asia*. Bangkok: Asia Books.
- Robson, C. R., Buck, H., Farrow, D. S., Fisher, T. and King, B. F. (1998) A birdwatching visit to the Chin Hills, West Burma (Myanmar), with notes from nearby area. *Forktail* 13: 109-115.
- Round, P. D. (1988) *Resident forest birds in Thailand: their status and conservation*. Cambridge, U.K.: International Council for Bird Preservation.
- Tordoff, A., Swan, S., Grindley, M. and Siurua, H. (1999) *Hoang Lien Nature Reserve: biodiversity survey and conservation evaluation 1997/8*. Frontier-Vietnam Forest Research Programme Technical Report No. 13. London and Hanoi: the Society for Environmental Exploration and the Institute of Ecology and Biological Resources, Hanoi.
- Vo Quy and Nguyen Cu (1995) [*Checklist of the birds of Vietnam*]. Hanoi: Agricultural Publishing House. (In Vietnamese.)

Andrew W. Tordoff, Society for Environmental Exploration, 77 Leonard St, London, EC2A 4QS, U.K. Correspondence and present address: BirdLife International Vietnam Programme, 11 Lane 167, Tay Son, Hanoi, Vietnam. Email: jack@birdlife.netnam.vn

White Tern *Gygis alba* sighted at Narcondam Island, Bay of Bengal, India

H.S.A. YAHYA and ASHFAQ AHMAD

Narcondam Island (13°25'N 94°17'E) lies in the Bay of Bengal between the Andaman Islands and the coast of Burma (Hussain 1984). The island is a part of a submerged line of peaks, which include the Andaman and Nicobar Islands, and lies 79 miles north-east of Port Blair in the Andaman islands (Abdulali 1971). Its

area is approximately 6.8 km², and the highest point reaches about 530 m above sea level.

We conducted a short study during March 2000 to assess the status of the endemic Narcondam Hornbill *Rhyticeros narcondami*. During the study we also prepared a bird checklist for comparison with previous records.